

Would you like your staff to be expert on FASD?

Is it safe to drink alcohol while pregnant?

Alcohol is a substance that is known to cause birth defects. Other factors can increase or decrease the potential of alcohol to cause harm to the fetus but there are so many factors that no one can say for sure that it is safe to drink some alcohol. The rffada recommends that for a healthy baby, no alcohol is the safest choice¹.

What are these 'other factors'?

Factors apart from alcohol that may play a role in the severity of the condition include:

- Nutrition
- Other drug use
- Stress
- Epigenetics | genetics
- Number of births | pregnancies
- Height | weight

These are some reasons why prenatal exposure to alcohol may affect one baby more severely than another.

It is impossible to tell how even one unit of alcohol will affect your baby so the rffada advises that no alcohol while pregnant, breastfeeding or trying to become pregnant is the safest option.

<http://sogc.org/guidelines/alcohol-use-and-pregnancy-consensus-clinical-guidelines/>

The Russell Family Fetal Alcohol Disorders Association (*rffada*) is an unfunded not-for-profit health promotion charity dedicated to ensuring that individuals affected prenatally by alcohol have access to services and support

Russell Family



Fetal Alcohol Disorders ASSOCIATION INC

Elizabeth (Anne) Russell
PO Box 6795
CAIRNS Q 4870
M - 0412 550 540
W - www.rffada.org
E - elizabeth@rffada.org



training outside the box

For training on Fetal Alcohol Spectrum Disorder

Contact Training Connections Australia
0 4 1 2 5 5 0 5 4 0



No safe time and no safe amount

Russell Family Fetal Alcohol Disorders ASSOCIATION INC



Prepared as a community service by the
Russell Family Fetal Alcohol Disorders
Association and sponsored by -



ALCOHOL CAN HARM AN UNBORN BABY

What is Fetal Alcohol Spectrum Disorder?

What is FASD? Is it the new thalidomide?

Fetal Alcohol Spectrum Disorder (FASD) is an umbrella term used to describe a range of disabilities that can be caused through prenatal exposure to alcohol.

FASD is widely recognised as the most common preventable cause of birth defects and brain damage in the world.

FASD is not a diagnosis. It represents a range of possible diagnoses that fall under the spectrum. These diagnoses are Fetal Alcohol Syndrome (FAS), partial Fetal Alcohol Syndrome (pFAS), Alcohol Related Neuro-developmental Disorders (ARND) and Alcohol Related Birth Defects (ARBD).

If doctors have not been specially trained to diagnose FASD, they may misdiagnose affected children with ADD |ADHD, Autism | Asperger's, ODD, GDD, RAD, and other similar conditions. Experts say that the more of these diagnoses a child has, the more likely it is to be FASD.

At right: Seth Russell
24 months

Seth is now 31 years old. He has full Fetal Alcohol Syndrome



But, what does all that mean?

If a child has FASD, he or she may have some of the problems below. However with an early diagnosis and appropriate interventions, some of these problems can be reduced. If you think your child has FASD, talk to your doctor or contact Anne at elizabeth@rffada.org

- Affected babies may have trouble settling
- May be hyperactive and difficult to parent
- Difficulties at school particularly in the higher grades
- Difficulty linking cause and effect
- 'Normal' parenting techniques do not work such as natural justice or tough love
- Inability to generalise learning from one place to another
- Memory may work one day but not the next
- Most affected people will have a normal IQ
- Very likely to have a mental illness and drug or alcohol addiction
- Difficulty keeping friends or relationships
- May experience trouble with the police
- Problems maintaining employment
- Inappropriate sexual behaviour
- Prison |incarceration | mental health facility

Website	www.rffada.org
Like	Russell Family Fetal Alcohol Disorders Assn
Like	rffada – parents and carers group
Like	rffada – living with FASD
Like	rffada – WA FASD Support Group
Like	Queensland FASD Support Group

I had a drink before I knew I was pregnant - what should I do?

If you have been drinking small amounts do **not** worry – however the sooner you stop drinking the **healthier** your baby will be. Talk to your doctor or contact Anne at elizabeth@rffada.org.

As you can see in the information in this brochure, there are many factors can make a positive or negative difference to the health of your baby.

Where can I find out more?

Visit www.rffada.org for information on FASD or email elizabeth@rffada.org.

Australia-wide training on FASD is available. We can train Employment Consultants; Police; Corrections; Teachers; Foster carers; Alcohol and drug workers; Parents and carers; Frontline workers; Mental health workers and others.



At left: Mick Russell on his first day at Kindergarten. Mick has Alcohol Related Neuro-developmental Disorder.

Mick is now 34 years of age.



NO alcohol is the safest choice for baby